

Title	Liver-specific γ -glutamyl carboxylase-deficient mice display bleeding diathesis and short life span
Author(s)	Azuma, Kotaro; Tsukui, Tohru; Ikeda, Kazuhiro et al.
Citation	PLoS ONE. 9(2) p.e88643
Issue Date	2014-02-10
oaire:version	VoR
URL	https://hdl.handle.net/11094/78602
rights	© 2014 Azuma et al. This article is licensed under a Creative Commons Attribution 4.0 International License.
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Liver-Specific γ -Glutamyl Carboxylase-Deficient Mice Display Bleeding Diathesis and Short Life Span

Kotaro Azuma¹, Tohru Tsukui², Kazuhiro Ikeda², Sachiko Shiba², Kimie Nakagawa³, Toshio Okano³, Tomohiko Urano⁴, Kuniko Horie-Inoue², Yasuyoshi Ouchi¹, Masahito Ikawa⁵, Satoshi Inoue^{1,2,4*}

1 Department of Geriatric Medicine, Graduate School of Medicine, The University of Tokyo, Bunkyo-ku, Tokyo, Japan, **2** Division of Gene Regulation and Signal Transduction, Research Center for Genomic Medicine, Saitama Medical University, Hidaka, Saitama, Japan, **3** Department of Hygienic Sciences, Kobe Pharmaceutical University, Kobe, Hyogo, Japan, **4** Department of Anti-Aging Medicine, Graduate School of Medicine, The University of Tokyo, Bunkyo-ku, Tokyo, Japan, **5** Research Institute for Microbial Diseases, Osaka University, Suita, Osaka, Japan

Abstract

Vitamin K is a fat-soluble vitamin that plays important roles in blood coagulation and bone metabolism. One of its functions is as a co-factor for γ -glutamyl carboxylase (Ggcx). Conventional knockout of *Ggcx* causes death shortly after birth in homozygous mice. We created *Ggcx*-floxed mice by inserting *loxP* sequences at the sites flanking exon 6 of *Ggcx*. By mating these mice with albumin-Cre mice, we generated *Ggcx*-deficient mice specifically in hepatocytes (*Ggcx* ^{Δ liver/ Δ liver} mice). In contrast to conventional *Ggcx* knockout mice, *Ggcx* ^{Δ liver/ Δ liver} mice had very low activity of Ggcx in the liver and survived several weeks after birth. Furthermore, compared with heterozygous mice (*Ggcx*^{+/ Δ liver}), *Ggcx* ^{Δ liver/ Δ liver} mice had shorter life spans. *Ggcx* ^{Δ liver/ Δ liver} mice displayed bleeding diathesis, which was accompanied by decreased activity of coagulation factors II and IX. *Ggcx*-floxed mice can prove useful in examining Ggcx functions *in vivo*.

Citation: Azuma K, Tsukui T, Ikeda K, Shiba S, Nakagawa K, et al. (2014) Liver-Specific γ -Glutamyl Carboxylase-Deficient Mice Display Bleeding Diathesis and Short Life Span. PLoS ONE 9(2): e88643. doi:10.1371/journal.pone.0088643

Editor: Peter J. Lenting, Institut National de la Santé et de la Recherche Médicale, France

Received: October 9, 2013; **Accepted:** January 6, 2014; **Published:** February 10, 2014

Copyright: © 2014 Azuma et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by grants from Cell Innovation Program, Grants-in-Aid, and Support Project of Strategic Research Center in Private Universities from the Ministry of Education, Culture, Sports, Science and Technology, Grants-in-Aid from the Ministry of Health, Labour, and Welfare, and the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: INOUE-GER@h.u-tokyo.ac.jp

Introduction

Vitamin K is a fat-soluble vitamin, which is involved in blood coagulation and bone metabolism. One of the major functions of vitamin K is its role as a co-factor for γ -glutamyl carboxylase (Ggcx) [1]. Ggcx is responsible for the posttranslational modification of glutamic acid (Glu) residues into γ -carboxyglutamic acid (Gla) by its carboxylase activity. Hitherto, 19 kinds of Gla proteins have been found, that is, coagulation factors II, VII, IX, and X; protein C, protein S, and protein Z [2]; osteocalcin [3]; nephrocalcin [4]; matrix Gla protein [5]; growth arrest specific-6 (Gas6) [6]; periostin [7]; β Ig-H3 [7]; proline-rich Gla protein 1 and 2 [8]; transmembrane Gla protein 3 and 4 [9]; upper zone of growth plate and cartilage matrix associated protein (UCMA; also called Gla-rich protein, GRP) [10]; and Ggcx itself, which was also shown to be γ -carboxylated [11]. Considering the various expression sites and functions of these Gla proteins, it is indicated that vitamin K is involved in many physiological and pathological processes by activating Ggcx.

On the other hand, we have previously demonstrated γ -carboxylation-independent vitamin K function, in which vitamin K is involved in the transcriptional regulation of nuclear receptor SXR/PXR [12]. We reported that SXR/PXR-dependent vitamin K functions are actually involved in the biological process in osteoblasts [13] and hepatocellular carcinoma cells [14].

To fully understand the function of vitamin K, it is vital to separate Ggcx-dependent and SXR/PXR-dependent vitamin K

functions. Analysis of *Ggcx* knockout mice would be useful in examining Ggcx-dependent vitamin K functions in each tissue; however, this attempt has been hampered by the fact that *Ggcx* knockout mice die between embryonic day 9.5 and 18, and the few that survive to term die shortly after birth [15]. To overcome this limitation, we generated *Ggcx*-floxed mice that enabled organ specific deficiency of Ggcx when bred with transgenic Cre mice that showed organ-specific expression of Cre recombinase. Here, we report a phenotype with liver-specific deficiency of Ggcx.

Materials And Methods

Ethics Statement

This study was carried out in strict accordance with the consent of the Animal Care and Use Committees of Osaka University and Saitama Medical University. The protocol was approved by the Committee on the Ethics of Animal Experiments of the University of Osaka (Protocol Number: 2003G004) and Saitama Medical University (Protocol Number: 797).

Targeting Vector Construction

A targeting vector was constructed using pNT1.1 containing two *loxP* sequences, a phosphoglycerate kinase (pgk)-neomycin selectable marker cassette (neomycin cassette), and a herpes simplex virus thymidine kinase gene. A mouse 129 strain λ genomic library (Stratagene) was purchased and digested with

NotI. Genomic fragments were subcloned into pBluescript SK(+). The 5' homology arm of the construct was derived from an Asp718/HindIII genomic fragment containing intron 4, exon 5, and intron 5 of *Ggcx*. This fragment was subcloned into pBluescript SK(+) and then inserted into the 5' region of pNT1.1 between the NotI and SalI sites. The 3' homology arm of the construct was derived from a genomic fragment containing intron 6, exon 7, and intron 7 of *Ggcx*. This fragment was amplified with primers 5'-GCTTAATTAATGCATATAAGACAAGCACC-3' and 5'-ATGGTACCTAGGAAAGCAGGAAGAAG-3' and inserted into the 3' region of pNT1.1 at the PacI and Asp718 sites. The genomic region containing exon 6 was amplified with primers 5'-AAGCTTGCAGGTGATTCTCC-3' and 5'-ATGCATAAACAGAAAAAGTGAGCAAGCC-3'; it was then inserted into pNT1.1 at a BamHI site between the 5'-*loxP* site and neomycin cassette. This resulted in a targeting vector with a neomycin cassette between exon 6 and 7 and a thymidine kinase gene located downstream of the 3' homology region. The targeting construct was linearized with NotI and electroporated into D3 ES cells [16].

Generation of *Ggcx*^{flax/flax} mice

Colonies of ES cells carrying the recombinant allele were screened using 150 µg/ml of G418 and negatively selected using 2 µM gancyclovir. Selected cells were amplified and genomic DNA was screened by Southern blot analysis. The ES cell lines carrying the recombinant allele were subsequently used to generate chimeras by injection into 129/Sv blastocysts. The chimeric mice were mated with wild type C57BL/6N mice. The F₁ agouti offspring were analyzed for homologous recombination by Southern blotting and PCR analysis. The F₁ offspring were backcrossed to C57BL/6N mice for more than eight generations to generate *Ggcx*^{flax/+} mice with a C57BL/6N genetic background. *Ggcx*^{flax/+} mice were intercrossed to generate *Ggcx*^{flax/flax} mice containing homozygous recombinant alleles.

Generation of hepatocyte-specific Ggcx-deficient mice

C57BL/6J mice containing transgenic constructs of mouse albumin enhancer/promoter and Cre recombinase modified to include a nuclear localization sequence (Alb-Cre) were purchased from the Jackson Laboratory. ROSA26-LacZ reporter mice were also obtained from the Jackson Laboratory. Hepatocyte-specific expression of Cre recombinase was confirmed by mating Alb-Cre mice with ROSA26-LacZ mice and assessing the β-galactosidase activity of the expressed *LacZ* gene, which is expected to be detected in cells expressing functional Cre recombinase. To generate hepatocyte-specific Ggcx-deficient mice (*Ggcx*^{Aliver/Aliver}), Alb-Cre mice were mated with *Ggcx*^{flax/flax} mice and F₁ offspring were subsequently intercrossed.

Southern blotting

EcoRI digested genomic DNA—derived from ES cells or tail specimens—was electrophoresed through a 0.6% agarose gel, transferred to a Hybond N+ membrane (Amersham Bioscience), and hybridized with the ³²P-labeled 164-bp sequence (ACAGCTTTCTTGATGCTGCTGGACATTCCCCAGGAAC-GCGGCCCTAGCTCCCTGGACCGAAAAATACTTGGATGG-GCTGGATGTGTGCCGTTTCCCCCTTGCTGGATGCCTT-GCGCCCACTGCCACTGGACTGGATGTATCTTGTCTAC-ACCATCATGTTTCTGGG) in exon 3 of the *Ggcx* gene.

Genotyping

Genomic DNA derived from tail specimens was used as the template for PCR analysis. Tail cut was done before 3 weeks old or immediately after the mice died. The Cre recombinase gene was detected by amplifying a 654-bp fragment within the *Cre* gene with primers 5'-CCTGGAAAATGCTTCTGTCCGTTTGCC-3' and 5'-GAGTTGATAGCTGGCTGGTGGCAGATG-3'. The *loxP* sequence was detected by amplifying the *Ggcx* sequence with primers 5'-AACTTAGGGAGTTGGTTCTCTTTCACTT-3' and 5'-AATCCAATACACCCAAGGTCTTATTCAT-3' in intron 5, containing *loxP* and linker sequences, to yield a 454-bp fragment from the *loxP*-containing allele and 407-bp fragment from the wild type allele. Deletion of exon 6 in the liver was confirmed with primers 5'-CGTGTACTTCATCGCGGGTG-3' within exon 6 and 5'-TCTGTATCCGGCTGAACGGG-3' within intron 6. DNA samples derived from liver, spleen, kidney and heart of both *Ggcx*^{Aliver/Aliver} mice and control *Ggcx*^{+/+} mice. The DNA samples of same concentration (3 ng/µl) were used as templates for PCR analysis.

Animal experiments

Mice were housed in a temperature-controlled room (22°C) with a 12-h light/dark schedule, had free access to water, and were fed standard laboratory chow. When mice were sacrificed, anesthesia with an intraperitoneal injection of 2.5% avertin was employed to minimize suffering of animals. Exsanguination was done following anesthesia to ensure death.

Ggcx activity assay

FLEEL was purchased from Bachem (Philadelphia, PA). α-phosphatidylcholine (type VE) and CHAPS were obtained from Sigma Aldrich Japan (Tokyo, Japan). Vitamin K₂ (menaquinone-4) was obtained from Eisai Co., Ltd. (Tokyo, Japan). The peptide ProFIX19, which contains the sequence AVFLDHENAN-KILNRPKRY, was synthesized by Genenet Co., Ltd. (Fukuoka, Japan). NaH¹⁴CO₃ (specific activity, 58 mCi/mmol) was obtained from Amersham Biosciences Corp. (NJ).

Six-week old mice were anesthetized with an intraperitoneal injection of 2.5% avertin and the livers were excised for measurement of Ggcx activity. Mice were euthanized by exsanguination following liver excision. The Ggcx activity was measured as previously described [17]. The amount of ¹⁴CO₂ incorporated into exogenous substrates was measured in reaction mixtures of 125 µl containing substrate (3.6 mM FLEEL), 222 µM reduced vitamin K (vitamin KH₂), 16 µM propeptide ProFIX19, 1.4 mM NaH¹⁴CO₃ (5 µCi), 25 mM MOPS (pH 7.0), 500 mM NaCl, 0.16% (w/v) phosphatidylcholine, 0.16% (w/v) CHAPS, 8 mM DTT, and 0.8 M ammonium sulfate, unless stated otherwise. All of the assay components, except for the microsomal fraction, were prepared as master mixes. ¹⁴CO₂ incorporation into peptide substrates (after an incubation period of over 30 min) was assayed using a scintillation counter. All assays were performed in quadruplicate.

Coagulation factor activity assay

Blood was collected from 6-week-old mice under anesthesia with an intraperitoneal injection of 2.5% avertin. Collected blood was immediately combined with one-tenth volume of 110 mM sodium citrate. Plasma was isolated by centrifugation for 15 min at 2500×g. The obtained plasma was analyzed with an automated blood coagulation analyzer (STA Compact, Roche, Basel, Switzerland) to determine factor II and IX activity using prothrombin (factor II) or factor IX-deficient plasma.

Bleeding test

Four-week-old mice were anesthetized with an intraperitoneal injection of 2.5% avertin. Their tails were cut to yield the same wound diameters. To evaluate bleeding time, filter paper was applied to the edge of the wound every minute, taking care not to dislodge the clot.

Hematological examination

Two ml of blood was collected from 6-week-old mice under anesthesia with an intraperitoneal injection of 2.5% avertin. Collected blood was mixed with anti-coagulants (1 mg of EDTA-2K and 20 μ l of 3% EDTA-3K). The number of platelets was measured using the Advia 120 (Bayer, Dublin, Ireland).

Life span analysis

To evaluate lifespan, mice were kept with their littermates. Male and female mice were kept in separate cages without mating. They were kept until either natural death, or evidence of impending mortality necessitating euthanasia, such as unresponsiveness to touch, slow respiration, coldness to touch, a hunched up position with matted fur. Condition of the mice was monitored every two days.

Statistical analysis

Data are expressed as mean \pm SEM. Differences between the mean values were analyzed using the unpaired Student's *t*-test. Survival rates were plotted using the Kaplan-Meier method. Survival differences between the groups were analyzed using the log-rank test, for which *p*-values were adjusted by the Bonferroni method.

Results

Generation of hepatocyte-specific *Ggcx*-deficient mice

The mouse γ -glutamyl carboxylase (*Ggcx*) gene consists of 15 exons (Figure 1A). To disrupt the *Ggcx* gene, the targeting vector was designed to flank exon 6 with two *loxP* sequences, and a frameshift was generated by excision with Cre recombinase (Figure 1A). Insertion of *loxP* sequences by homologous recombination was confirmed with Southern blotting analysis (Figure 1B). To delete the *Ggcx* gene in the liver alone, albumin-Cre (Alb-Cre) transgenic mice were used. The cre recombinase gene is under the control of the albumin promoter, which is active only in hepatocytes from E16.5 embryos [18] and the full activity was exhibited at 2 months after birth [19]. To confirm the specificity of recombination, the Alb-Cre mice were crossed with ROSA26-LacZ mice, which contain a reporter gene in which β -galactosidase is expressed in any tissue, and expression is dependent on Cre-mediated recombination. β -galactosidase activity was detected in the hepatocytes of offspring born from the mating of Alb-Cre with ROSA26-LacZ mice (Figure 2A); these offspring also showed Cre-recombinase activity in hepatocytes. Alb-Cre mice were then mated with *Ggcx*-floxed mice (*Ggcx^{lox/lox}*) and the resulting F₁ offspring were intercrossed. To examine the genotypes of the F₂ offspring, the Cre recombinase gene and the *loxP*-containing region of the *Ggcx* gene were amplified by PCR using genomic DNA prepared from tail samples. Some mice that expressed Cre recombinase and carried homozygous floxed alleles were considered to be liver-specific *Ggcx*-deficient mice (*Ggcx^{Δliver/Δliver}*) (Figure 2B). They were born alive and survived for at least several weeks. To confirm the ablation of *Ggcx* in the livers of *Ggcx^{Δliver/Δliver}* mice, genomic DNA was extracted from liver and other organs (spleen, kidney and heart) from 6-week old *Ggcx^{Δliver/Δliver}* mice and control *Ggcx^{+/+}* mice. Decreased intensity

of PCR products from exon 6 was observed in only livers of *Ggcx^{Δliver/Δliver}* mice (Figure 2C). Next, vitamin K-dependent *Ggcx* activity was measured in the livers of 6-week old *Ggcx^{Δliver/Δliver}* mice and control littermates (*Ggcx^{+/+}* mice). *Ggcx* activity was significantly decreased in the livers of *Ggcx^{Δliver/Δliver}* mice (Figure 2D). There was no significant difference in *Ggcx* activity between male and female *Ggcx^{Δliver/Δliver}* mice.

Bleeding diathesis in *Ggcx^{Δliver/Δliver}* mice

To examine the effect of decreased *Ggcx* activity in the livers of *Ggcx^{Δliver/Δliver}* mice, the activities of vitamin K-dependent coagulation factors were examined. Activities of factors II and IX were significantly decreased in *Ggcx^{Δliver/Δliver}* mice, compared with control (*Ggcx^{+/+}*) mice (Figure 3A). Decreased activity of vitamin K-dependent coagulation factor caused bleeding diathesis in *Ggcx^{Δliver/Δliver}* mice. Wild-type mice ceased bleeding within 10 minutes of tail incision, while *Ggcx^{Δliver/Δliver}* mice continued to bleed for more than 30 minutes (Figure 3B). The platelet count was not significantly different between wild type mice and *Ggcx^{Δliver/Δliver}* mice (Figure 3C), suggesting the longer bleeding time in *Ggcx^{Δliver/Δliver}* mice was due to defective secondary coagulation.

Shorter life span of *Ggcx^{Δliver/Δliver}* mice

As a result of bleeding diathesis, injury and pregnancy caused fatal bleeding in *Ggcx^{Δliver/Δliver}* mice. In 9-week-old *Ggcx^{Δliver/Δliver}* mice, massive subcutaneous bleeding was observed even before death. Dissection of pregnant *Ggcx^{Δliver/Δliver}* mice just after death revealed uterine as well as vaginal bleeding. Next, we evaluated the life span of *Ggcx^{Δliver/Δliver}* mice by keeping them separately without mating. Male *Ggcx^{Δliver/Δliver}* mice began to die from day 27 after birth, and all *Ggcx^{Δliver/Δliver}* male mice died within 80 days after birth (Figure 4). Female *Ggcx^{Δliver/Δliver}* mice began to die from day 39 after birth and 7 out of 11 (63.6%) survived longer than 100 days, unless they became pregnant (Figure 4). None of the control heterozygous littermates (*Ggcx^{+/Δliver}* mice) died within the 100 days of the observation period. The shorter life span of male *Ggcx^{Δliver/Δliver}* mice was statistically significant compared with male heterozygous littermates. The cause of death seemed to be anemia secondary to bleeding, since subcutaneous bleeding was observed in some *Ggcx^{Δliver/Δliver}* mice before death. Interestingly, female *Ggcx^{Δliver/Δliver}* mice survived significantly longer than male *Ggcx^{Δliver/Δliver}* mice.

Discussion

Mediation of post-transcriptional modification of substrate proteins by *Ggcx* is one of the major functions of vitamin K. So far, 19 proteins are known to be substrates of *Ggcx* and are expressed throughout body, indicating various physiological functions of vitamin K.

In the present study, we showed that liver-specific deficiency of *Ggcx* caused bleeding diathesis and short life span. We consider the massive bleeding in subcutaneous tissue or body cavity is a direct cause of death since we observed massive subcutaneous bleeding in most of the dead mice. It is also possible that local bleeding in vital organs such as brain can cause death due to bleeding diathesis.

Short life span of liver-specific *Ggcx*-deficient mice in the present study along with the clinical presentation of vitamin K deficiency indicate the relative importance of hepatic coagulation factors among *Ggcx* substrates. Coagulation factors II, VII, IX, and X are known to be vitamin K dependent. Therefore, we considered the decreased activity of these coagulation factors to be




Figure 1. Cre-recombinase-mediated tissue-specific excision of *Ggcx*. A, Strategy for homologous recombination. Targeting vector was designed to flank exon 6 of the *Ggcx* gene, and frame shift was generated by excision with Cre recombinase. Two *loxP* sequences (triangles) were inserted into introns 5 and 6. Neomycin cassette and *EcoRI* site were inserted into intron 6. B, Southern blot analysis of tail DNA. Homologous recombinant allele generated a 7.4-kb fragment by *EcoRI* digestion. A representative figure is shown.
doi:10.1371/journal.pone.0088643.g001

responsible for the *Ggcx*-deficient phenotype. Interestingly, although the activity of factors II and IX was decreased in *Ggcx*^{*Δliver/Δliver*} mice, they live much longer than those with a systemic lack of *Ggcx*. Most mice systemically lacking *Ggcx* die between embryonic day 9.5 and 18, and the few that survive to term die shortly after birth [15]. Among mice in which genes for vitamin K-dependent coagulation factors had been knocked out, factor II-deficient mice [20] and factor X-deficient mice [21] are partial embryonic lethal. In factor II-deficient fetuses, abnormal phenotypes such as pale yolk sac membrane, empty blood vessels, enlarged pericardial sacs, and distended hearts were observed, which appeared from embryonic day 9.5 to 12.5. In factor X-deficient mice, some fetuses began to die of massive bleeding from embryonic day 11.5 to 12.5, but the blood vessels and yolk sacs of these mice were normal. Considering the phenotypes of factor II-deficient and factor X-deficient mice, it can be inferred that the embryonic lethal phenotype of systemic *Ggcx*-deficient mice is likely due to abnormalities that developed at midgestation. In the present study, we used an albumin promoter to regulate Cre transcription. The albumin promoter is activated around embryonic day 16.5 [18]; therefore, *Ggcx* exists in the liver of *Ggcx*^{*Δliver/Δliver*} mice until embryonic day 16.5. This will contribute to a difference between liver-specific and systemic *Ggcx*-deficient mice, the latter lack *Ggcx* from the beginning of embryogenesis. Another possible reason that contributes to the difference between these *Ggcx*-deficient mice is incomplete ablation of *Ggcx* in *Ggcx*^{*Δliver/Δliver*} mice. We detected slight residual GGCX activity in the liver of *Ggcx*^{*Δliver/Δliver*} mice (Figure 2D) and factors II and IX activities were also detectable in the blood of *Ggcx*^{*Δliver/Δliver*} mice (Figure 3A). In another report using Alb-Cre mice, complete ablation of target gene was observed in 2 months after birth [19]. Thus, it is assumed that residual *Ggcx* activity can also remain for several weeks after birth. These slight residual activities may have been vital for the survival of *Ggcx*^{*Δliver/Δliver*} mice.

Both factor VII-deficient mice [22] and factor IX-deficient mice [23] displayed bleeding diathesis. The factor IX-deficient mice showed swollen extremities and extensive hemorrhagic lesions following mechanical trauma, although they survived for at least several weeks. In contrast, the factor VII-deficient mice survived to term and followed a normal Mendelian inheritance pattern. However, most of them died perinatally owing to intra-abdominal hemorrhage within 24 hours, and the remaining neonates died from intracranial hemorrhage in 24 days. Considering the aggressive bleeding of factor VII-deficient mice, the residual activity of *Ggcx* in *Ggcx*^{*Δliver/Δliver*} mice may contribute to the survival. Furthermore, *Ggcx* activity before embryonic day 16.5 may have some preventive effect against postnatal bleeding.

In regard to the phenotypes of conditional deficiency of coagulation factors, factor VII-insufficient mice at the 0.7% expression level compared with wild-type mice could survive to adulthood despite displaying severely downregulated overall thrombin production and cardiac fibrosis at a young adult age [24]. Induction of prothrombin ablation in adulthood using Mx1-Cre caused fatal hemorrhagic events particularly in heart and brain [25]. Liver-specific *Ggcx*-deficient mice in the present study exhibit a longer life span in comparison with that of prothrombin deletion in adult mice, because the amount of coagulation factors in *Ggcx*-deficient mice are substantially decreased but even sufficient to survive for several weeks after birth. In comparison with factor VII-insufficient mice, however, it is assumed that severe insufficiency of multiple coagulation factors occurred in liver-specific *Ggcx*-deficient mice simultaneously.

It is intriguing that mice lacking fibrinogen, the final effector of the coagulation cascade, displayed similar phenotypes to those seen in *Ggcx*^{*Δliver/Δliver*} mice [26]. They suffered from spontaneous abdominal hemorrhage, but long term survival was possible. In fibrinogen-deficient mice, pregnant female ones died from vaginal hemorrhage, which was also observed in *Ggcx*^{*Δliver/Δliver*} mice.




Figure 2. Liver-specific ablation of *Ggcx* gene. A, The albumin promoter is active only in hepatocytes. Alb-Cre or wild type (WT) mice were mated with ROSA26-LacZ mice. Livers were obtained postnatally from mice and stained with X-gal. β -galactosidase activity was detected in the hepatocytes of the mouse born from mating of Alb-Cre and ROSA26-LacZ mice (right panel). Liver of the mouse born from wild type was shown as a negative control (left panel). Representative figures are shown for each group. B, Genotyping using genomic tail DNA. Expected bands from Cre recombinase, *loxP* sequence (floxed allele), and wild type allele are shown. In mouse #2, #4 and #5, both alleles were replaced with *loxP* containing recombinant alleles, with at least one copy of Cre recombinase. A representative result is shown. C, Liver-specific ablation of *Ggcx* gene was confirmed with PCR analysis. DNA samples derived from liver, spleen, kidney and heart of 6-month week old *Ggcx*^{Δliver/Δliver} mice (4 male mice and 4 female mice) and control *Ggcx*^{+/+} mice (4 male mice and 4 female mice) were used as templates. D, Activity of *Ggcx* in the liver-specific *Ggcx*-deficient mice. Microsome was prepared from the livers of 6-week old *Ggcx*^{Δliver/Δliver} mice and control *Ggcx*^{+/+} mice of both sexes. The activity of *Ggcx* was measured by ¹⁴CO₂ incorporated into exogenous substrate in the presence of reduced vitamin K (222 μ M). Bars represent the mean value \pm SEM (n = 4). Differences between the mean values were analyzed using the unpaired Student's *t*-test. ***P* < 0.01. doi:10.1371/journal.pone.0088643.g002

In this study, we observed longer life spans of female *Ggcx*^{Δliver/Δliver} mice compared with male *Ggcx*^{Δliver/Δliver} mice. Notably, this sexual dimorphism of life span was also observed in fibrinogen-deficient mice [26], although the difference was smaller than that of *Ggcx*^{Δliver/Δliver} mice. Considering activities of *Ggcx* in the livers of *Ggcx*^{Δliver/Δliver} mice were not significantly different between male and female (Figure 2D), this sexual dimorphism may be owing to the difference in aggressiveness of behavior between males and females. Typically, males are more aggressive than females [27], which causing males more susceptible to injury. Another explanation for the sexual dimorphism of

life span is the procoagulant activity of female sex hormone, estrogen. Clinically estrogen administration as oral contraceptives or hormone replacement therapy is known to be associated with higher risk of venous thrombosis [28]. In experiment using rats, estrogen was shown to prevent decline of prothrombin caused by vitamin K deficiency [29]. This report suggests higher concentration of estrogen in female mice may ameliorate bleeding diathesis due to *Ggcx* deficiency which is pathologically similar with vitamin K deficiency.

Interestingly, we observed higher activities of *Ggcx* as well as vitamin K-dependent coagulation factors in wild type male mice






Figure 3. Bleeding diathesis in *Ggcx*^{Δliver/Δliver} mice. A, Decreased activity of coagulation factor in *Ggcx*^{Δliver/Δliver} mice. Activities of factors II and IX were significantly decreased in *Ggcx*^{Δliver/Δliver} mice compared with wild type (*Ggcx*^{+/+}) mice. Bars represent the mean value \pm SEM (n=8). Differences between the mean values were analyzed using the unpaired Student's *t*-test. ****P*<0.001. B, Prolonged bleeding time in *Ggcx*^{Δliver/Δliver} mice. Tail bleeding time was measured by a filter paper method. *Ggcx*^{Δliver/Δliver} mice continued bleeding for more than 30 min. Gray triangle: 0 min; Black triangle: 10 min. A representative figure is shown. C, Platelet counts of *Ggcx*^{Δliver/Δliver} mice. Hematological examination of male *Ggcx*^{Δliver/Δliver} mice (n=6), female *Ggcx*^{Δliver/Δliver} mice (n=12), male wild type mice (n=10), and female wild type mice (n=10) was performed. The platelet count was not significantly different between wild type mice and *Ggcx*^{Δliver/Δliver} mice. Bars represent the mean value \pm SEM. NS: not significant. doi:10.1371/journal.pone.0088643.g003

than in wild type female mice (Figure 2D and 3A). In the study by Wong et al. [30], some of the procoagulant factor activity levels were higher in male mice than in female mice and were growth hormone-dependent. In the study by Kundu et al., factor IX




Figure 4. Shorter life span of *Ggcx*^{Δliver/Δliver} mice. Cumulative life spans of male *Ggcx*^{Δliver/Δliver} mice (n=10), female *Ggcx*^{Δliver/Δliver} mice (n=11), male heterozygous littermates (*Ggcx*^{+/-} mice) (n=12), and female heterozygous littermates (*Ggcx*^{+/-} mice) (n=12) were calculated by the Kaplan-Meier method and compared using the log-rank test. P-values were adjusted by the Bonferroni method. Life span of male *Ggcx*^{Δliver/Δliver} mice was significantly shorter (*P*<0.001) compared with that of heterozygous mice. The life spans of female *Ggcx*^{Δliver/Δliver} mice were significantly longer than those of male *Ggcx*^{Δliver/Δliver} mice (*P*=0.044). doi:10.1371/journal.pone.0088643.g004

activity might be higher in one-month-aged male mice compared with one-month-aged females [31]. Since growth hormone has been known to contribute to sexual dimorphism in liver protein expression [32], it is assumed that the hormone will also exert difference in *Ggcx* activity.

In human, two missense mutations of *GGCX* gene were reported to cause hereditary bleeding disorder due to low activity of vitamin K-dependent coagulation factors [33,34]. In our study about 60% of female *Ggcx*^{Δliver/Δliver} mice survived longer than 100 days indicated small portion of residual *Ggcx* activity detected in the liver of *Ggcx*^{Δliver/Δliver} mice is sufficient to survive unless they got injured or pregnant. This is compatible with the clinical observation that patients with decreased carboxylase activity live several years before diagnosis [33,34].

In the current study, we successfully generated mice exhibiting liver-specific insufficiency of *Ggcx* activity. Because systemic *Ggcx* knockout mice do not live long after birth, our animal model enables to show the phenotype of liver-specific *Ggcx* deficiency for the first time and also will open up the possibility to evaluate in extra-hepatic organ-specific *Ggcx* activities using Cre recombinase driven by proper organ-specific promoters. Recently, several clinical and epidemiological studies have suggested extrahepatic actions of vitamin K. Its fracture-prevention effect has been proven in clinical studies [35,36], and vitamin K2 is used as a drug for osteoporosis treatment in several Asian countries. In epidemiological studies, low serum concentration of vitamin K was reported to be correlated with osteoarthritis [37], dementia [38], and coronary artery disease [39,40]. Moreover, a biosynthetic enzyme of menaquinone, which is an active form of vitamin K, was found to be expressed in extrahepatic organs [41]. These discoveries along with the many *Ggcx* substrates expressed throughout the body, suggest the extrahepatic function of *Ggcx* is worth investigating. The *Ggcx*-floxed mice we created in this study would be useful in clarifying vitamin K action in the whole body.

Acknowledgments

We are grateful to A. Kawai, M. Tanaka, Y Esaki, and T. Tanaka for their technical assistance.

Author Contributions

Conceived and designed the experiments: KA TT SI. Performed the experiments: KA TT KI SS KN MI. Analyzed the data: KA TT KI SS

KN TO MI SI. Contributed reagents/materials/analysis tools: TO TU KHI YO MI SI. Wrote the paper: KA TT KI KHI SI.

References

- Furie B, Bouchard BA, Furie BC (1999) Vitamin K-dependent biosynthesis of gamma-carboxyglutamic acid. *Blood* 93: 1798–1808.
- Rezaie AR, Bac JS, Manithody C, Qureshi SH, Yang L (2008) Protein Z-dependent protease inhibitor binds to the C-terminal domain of protein Z. *J Biol Chem* 283: 19922–19926.
- Price PA, Otsuka AA, Poser JW, Kristaponis J, Raman N (1976) Characterization of a gamma-carboxyglutamic acid-containing protein from bone. *Proc Natl Acad Sci U S A* 73: 1447–1451.
- Worcester EM, Sebastian JL, Hiatt JG, Beshensky AM, Sadowski JA (1993) The effect of warfarin on urine calcium oxalate crystal growth inhibition and urinary excretion of calcium and nephrocalcin. *Calcif Tissue Int* 53: 242–248.
- Luo G, Ducey P, McKee MD, Pincro GJ, Loyer E, et al. (1997) Spontaneous calcification of arteries and cartilage in mice lacking matrix GLA protein. *Nature* 386: 78–81.
- Varnum BC, Young C, Elliott G, Garcia A, Bartley TD, et al. (1995) Axl receptor tyrosine kinase stimulated by the vitamin K-dependent protein encoded by growth-arrest-specific gene 6. *Nature* 373: 623–626.
- Coutu DL, Wu JH, Monette A, Rivard GE, Blostein MD, et al. (2008) Periostin, a member of a novel family of vitamin K-dependent proteins, is expressed by mesenchymal stromal cells. *J Biol Chem* 283: 17991–18001.
- Kulman JD, Harris JE, Haldeman BA, Davie EW (1997) Primary structure and tissue distribution of two novel proline-rich gamma-carboxyglutamic acid proteins. *Proc Natl Acad Sci U S A* 94: 9058–9062.
- Kulman JD, Harris JE, Xie L, Davie EW (2001) Identification of two novel transmembrane gamma-carboxyglutamic acid proteins expressed broadly in fetal and adult tissues. *Proc Natl Acad Sci U S A* 98: 1370–1375.
- Viegas CS, Simes DC, Laizé V, Williamson MK, Price PA, et al. (2008) Gla-rich protein (GRP), a new vitamin K-dependent protein identified from sturgeon cartilage and highly conserved in vertebrates. *J Biol Chem* 283: 36655–36664.
- Berkner KL, Pudota BN (1998) Vitamin K-dependent carboxylation of the carboxylase. *Proc Natl Acad Sci U S A* 95: 466–471.
- Tabb MM, Sun A, Zhou C, Grün F, Errandi J, et al. (2003) Vitamin K2 regulation of bone homeostasis is mediated by the steroid and xenobiotic receptor SXR. *J Biol Chem* 278: 43919–43927.
- Ichikawa T, Horie-Inoue K, Ikeda K, Blumberg B, Inoue S (2006) Steroid and xenobiotic receptor SXR mediates vitamin K2-activated transcription of extracellular matrix-related genes and collagen accumulation in osteoblastic cells. *J Biol Chem* 281: 16927–16934.
- Azuma K, Urano T, Ouchi Y, Inoue S (2009) Vitamin K2 suppresses proliferation and motility of hepatocellular carcinoma cells by activating steroid and xenobiotic receptor. *Endocr J* 56: 843–849.
- Zhu A, Sun H, Raymond RM Jr, Furie BC, Furie B, et al. (2007) Fatal hemorrhage in mice lacking gamma-glutamyl carboxylase. *Blood* 109: 5270–5275.
- Doetschman TC, Eistetter H, Katz M, Schmidt W, Kemler R (1985) The in vitro development of blastocyst-derived embryonic stem cell lines: formation of visceral yolk sac, blood islands and myocardium. *J Embryol Exp Morphol* 87: 27–45.
- Kinoshita H, Nakagawa K, Narusawa K, Goski-Sone M, Fukushi-Irie M, et al. (2007) A functional single nucleotide polymorphism in the vitamin-K-dependent gamma-glutamyl carboxylase gene (Arg325Gln) is associated with bone mineral density in elderly Japanese women. *Bone* 40: 451–456.
- Wan YJ, An D, Cai Y, Repa JJ, Hung-Po Chen T, et al. (2000) Hepatocyte-specific mutation establishes retinoid X receptor alpha as a heterodimeric integrator of multiple physiological processes in the liver. *Mol Cell Biol* 20: 4436–4444.
- Gu J, Weng Y, Zhang QY, Cui H, Behr M, et al. (2003) Liver-specific deletion of the NADPH-cytochrome P450 reductase gene: impact on plasma cholesterol homeostasis and the function and regulation of microsomal cytochrome P450 and heme oxygenase. *J Biol Chem* 278: 25895–25901.
- Sun WY, Witte DP, Degen JL, Colbert MC, Burkart MC, et al. (1998) Prothrombin deficiency results in embryonic and neonatal lethality in mice. *Proc Natl Acad Sci U S A* 95: 7597–7602.
- Dewerchin M, Liang Z, Moons L, Carmeliet P, Castellino FJ, et al. (2000) Blood coagulation factor X deficiency causes partial embryonic lethality and fatal neonatal bleeding in mice. *Thromb Haemost* 83: 185–190.
- Rosen ED, Chan JC, Idusogie E, Clotman F, Vlasuk G, et al. (1997) Mice lacking factor VII develop normally but suffer fatal perinatal bleeding. *Nature* 390: 290–294.
- Wang L, Zoppè M, Hackeng TM, Griffin JH, Lee KF, et al. (1997) A factor IX-deficient mouse model for hemophilia B gene therapy. *Proc Natl Acad Sci U S A* 94: 11563–11566.
- Rosen ED, Xu H, Liang Z, Martin JA, Suckow M, et al. (2005) Generation of genetically-altered mice producing very low levels of coagulation factor VII. *Thromb Haemost* 94: 493–497.
- Mullins ES, Kombrinck KW, Talmage KE, Shaw MA, Witte DP, et al. (2009) Genetic elimination of prothrombin in adult mice is not compatible with survival and results in spontaneous hemorrhagic events in both heart and brain. *Blood* 113: 696–704.
- Suh TT, Holmbäck K, Jensen NJ, Daugherty CC, Small K, et al. (1995) Resolution of spontaneous bleeding events but failure of pregnancy in fibrinogen-deficient mice. *Genes Dev* 9: 2020–2033.
- Gatewood JD, Wills A, Shetty S, Xu J, Arnold AP, et al. (2006) Sex chromosome complement and gonadal sex influence aggressive and parental behaviors in mice. *J Neurosci* 26: 2335–2342.
- Rosendaal FR, Helmerhorst FM, Vandenbroucke JP (2002) Female hormones and thrombosis. *Arterioscler Thromb Vasc Biol* 22: 201–210.
- Matschiner JT, Willingham AK (1974) Influence of sex hormones on vitamin K deficiency and epoxidation of vitamin K in the rat. *J Nutr* 104: 660–665.
- Wong JH, Dukes J, Levy RE, Sos B, Mason SE, et al. (2008) Sex differences in thrombosis in mice are mediated by sex-specific growth hormone secretion patterns. *J Clin Invest* 118: 2967–2978.
- Kundu RK, Sangiorgi F, Wu LY, Kurachi K, Anderson WF, et al. (1998) Targeted inactivation of the coagulation factor IX gene causes hemophilia B in mice. *Blood* 92: 168–174.
- Waxman DJ, O'Connor C (2006) Growth hormone regulation of sex-dependent liver gene expression. *Mol Endocrinol* 20: 2613–2629.
- Brenner B, Sánchez-Vega B, Wu SM, Lanir N, Stafford DW, et al. (1998) A missense mutation in gamma-glutamyl carboxylase gene causes combined deficiency of all vitamin K-dependent blood coagulation factors. *Blood* 92: 4554–4559.
- Spronk HM, Farah RA, Buchanan GR, Vermeer C, Soute BA (2000) Novel mutation in the gamma-glutamyl carboxylase gene resulting in congenital combined deficiency of all vitamin K-dependent blood coagulation factors. *Blood* 96: 3650–3652.
- Shiraki M, Shiraki Y, Aoki C, Miura M (2000) Vitamin K2 (menatetrenone) effectively prevents fractures and sustains lumbar bone mineral density in osteoporosis. *J Bone Miner Res* 15: 515–521.
- Cockayne S, Adamson J, Lanham-New S, Shearer MJ, Gilbody S, et al. (2006) Vitamin K and the prevention of fractures: systematic review and meta-analysis of randomized controlled trials. *Arch Intern Med* 166: 1256–1261.
- Neogi T, Booth SL, Zhang YQ, Jacques PF, Terkeltaub R, et al. (2006) Low vitamin K status is associated with osteoarthritis in the hand and knee. *Arthritis Rheum* 54: 1255–1261.
- Presse N, Shatenstein B, Kergoat MJ, Ferland G (2008) Low vitamin K intakes in community-dwelling elders at an early stage of Alzheimer's disease. *J Am Diet Assoc* 108: 2095–2099.
- Geleijnse JM, Vermeer C, Grobbee DE, Schurgers IJ, Knapen MH, et al. (2004) Dietary intake of menaquinone is associated with a reduced risk of coronary heart disease: the Rotterdam Study. *J Nutr* 134: 3100–3105.
- Gast GC, de Roos NM, Shuijs I, Bots ML, Beulens JW, et al. (2009) A high menaquinone intake reduces the incidence of coronary heart disease. *Nutr Metab Cardiovasc Dis* 19: 504–510.
- Nakagawa K, Hirota Y, Sawada N, Yuge N, Watanabe M, et al. (2010) Identification of UBIAD1 as a novel human menaquinone-4 biosynthetic enzyme. *Nature* 468: 117–121.